Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

TURKEY.

Report from Trebizond—Suspect plague.

Consul Jewett reports, August 9, as follows: Plague suspected.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

CANADA—*Quebec—Sherbrooke.*—Month of July, 1906. Estimated population, 13,950. Total number of deaths, 19, including 1 from enteric fever.

FRANCE—*Roubaix.*—Month of July, 1906. Estimated population, 124,660. Total number of deaths, 171, including diphtheria 1, measles 5, and 1 from whooping cough.

St. Etienne.—Two weeks ended July 15, 1906. Estimated population, 146,836. Total number of deaths, 109, including diphtheria 1, measles 1, whooping cough 1, and 21 from tuberculosis.

GERMANY—*Strasburg.*—Month of June, 1906. Estimated population, 180,000. Total number of deaths, 240, including diphtheria 3, enteric fever 3, measles 2, scarlet fever 2, and 32 from tuberculosis.

GIBRALTAR.—Four weeks ended July 29, 1906. Estimated population, 27,460. Total number of deaths, 31, including 3 from enteric fever.

GREAT BRITAIN—*England and Wales.*—The deaths registered in 76 great towns in England and Wales during the week ended July 28, 1906, correspond to an annual rate of 12.6 per 1,000 of population, which is estimated at 15,818,360.

Bradford.—Two weeks ended July 28, 1906. Estimated population, 288,544. Total number of deaths 135, including enteric fever 1, measles 4, scarlet fever 1, whooping cough 1, and 9 from phthisis pulmonalis.

London.—One thousand and fifty-six deaths were registered during the week, including measles 30, scarlet fever 6, diphtheria 10, enteric fever 3, whooping cough 10, and 74 from diarrhea. The deaths from all causes correspond to an annual rate of 11.7 per 1,000. In Greater London 1,476 deaths were registered. In the "outer ring" the deaths included 4 from diphtheria, 5 from measles, 1 from scarlet fever, and 2 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended July 28, 1906, in the 21 principal town districts of Ireland was 17.3 per 1,000 of the population, which is estimated at 1,093,959. The lowest rate was recorded in Dundalk, viz, 4, and the highest in Sligo, viz, 33.6 per 1,000. In Dublin and suburbs 149 deaths were registered, including measles 2, scarlet fever 2, whooping cough 2, and 32 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended July 28, 1906, correspond to an annual rate of 14.3 per

1,000 of the population, which is estimated at 1,787,788. The lowest rate of mortality was recorded in Perth, viz, 6, and the highest in Paisley, viz, 15 per 1,000. The aggregate number of deaths registered from all causes was 489, including diphtheria 4, enteric fever 9, measles 6, scarlet fever 2, and 11 from whooping cough.

JAMAICA—*Port Antonio.*—Two weeks ended June 30, 1906. Estimated population, 3,000. Number of deaths not reported. No deaths from contagious diseases.

JAPAN—*Formosa.*—Month of June, 1906. Estimated population, 3,050,004. Number of deaths not reported. Ten deaths from enteric fever and 514 from plague reported.

Ten days ended July 10, 1906, 66 deaths from plague reported.

JAVA—*Batavia.*—Two weeks ended June 30, 1906. Estimated population, 160,000. Total number of deaths not reported. No contagious diseases.

MALTA.—Weeks ended June 16 and 30, 1906. Estimated population, 205,059. Total number of deaths, 89 and 105, including diphtheria 1, and 2 from whooping cough.

Three weeks ended July 21, 1906. Total number of deaths, 379, including enteric fever 1, and 2 from whooping cough.

WEST INDIES—*Antigua.*—Three weeks ended July 21, 1906. Estimated population, 36,700. Total number of deaths, 24. No contagious diseases reported.

Curaçao.—Two weeks ended August 4, 1906. Estimated population, 31,600. Total number of deaths, 22. No contagious diseases.

Cholera, yellow fever, plague, and smallpox, from June 29 to August 24, 1906.

[Reports received by the Surgeon-General, Public Health and Marine-Hospital Service, from American consuls through the Department of State, and from other sources.]

[For reports received from December 29, 1905, to June 29, 1906, see PUBLIC HEALTH REPORTS for June 29, 1906.]

[NOTE.—In accordance with custom, the tables of epidemic diseases are terminated semiannually and new tables begun.]

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
China:				
Hongkong	July 7	1	1	
India:				
Bombay	May 30–Aug. 10		174	
Calcutta	May 20–July 7		214	
Karachi	July 2–8	1	1	
Madras	June 2–July 13		45	
Rangoon	May 27–June 30		9	
Philippine Islands:				
Manila	May 13–June 30	70	63	
Provinces	Apr. 28–June 30	262	159	1 on schr. Florence S., 1 on ss. Ban Yek, and 1 on ss. Vasco Navarro, at Mariveles.
Siam:				
Bangkok	Feb. 3–Apr. 14	224	198	
Straits Settlements:				
Parak	May 9–12	64	34	
Singapore	May 9–June 29	50	45	
Wellesley Province	Apr. 10–May 12	107	81	